

CP phase of nonuniversal Z' on $\sin \phi_s^{J/\Psi\phi}$ and T-odd observables of $\bar{B}_q \rightarrow V_q \ell^+ \ell^-$

Chuan-Hung Chen^{1,2*}

¹*Department of Physics, National Cheng-Kung University, Tainan 701, Taiwan*

²*National Center for Theoretical Sciences, Hsinchu 300, Taiwan*

(Dated: December 25, 2009)

Abstract

The evidence of a large CP phase has been shown by CDF and DØ collaborations in the time-dependent CP asymmetry (CPA) of $B_s \rightarrow J/\Psi\phi$ decay, where the nonvanished CPA clearly implies the existence of a non-Kobayashi-Maskawa phase in $b \rightarrow s$ transition. We study the new phases originated by the extra family dependent $U(1)'$ model and examine their impact on $\hat{T}$ or CP violating observables in $\bar{B}_q \rightarrow V_q \ell^+ \ell^-$ decays with $V_q = K^* (\phi)$. Adopting the constraints from the current data of Δm_s and the branching ratios for $\bar{B}_s \rightarrow \mu^+ \mu^-$ and $\bar{B}_q \rightarrow V_q \ell^+ \ell^-$, we find that $\phi_s^{J/\Psi\phi} = -(0.26 \pm 0.22)$ and the T-odd observables of above 10% in the decay chain $\bar{B}_q \rightarrow V_q (\rightarrow K\pi[KK]) \ell^+ \ell^-$ can be achieved. In addition, we demonstrate that the longitudinal and transverse polarizations of K^* and the up-down asymmetry of K in the same decay chain are also sensitive to the Z' -mediated effects.

* Email: physchen@mail.ncku.edu.tw

Although three generations of the standard model (SM) provide a unique CP violating phase in the Cabibbo-Kobayashi-Maskawa (CKM) matrix [1] to interpret the observed CP violating phenomena in K and B systems, due to the failure of the KM phase in the explanation of matter-antimatter asymmetry, some new source of CP violation (CPV) is expected. Therefore, it is an important issue to probe and find the new CP violating effects. By the B_s production in Tevatron Run II, CDF and DØ have observed the B_s oscillation of $\Delta m_s = 17.77 \pm 0.10 \pm 0.07 \text{ ps}^{-1}$ [3] and $\Delta m_s = 18.56 \pm 0.87 \text{ ps}^{-1}$ [4], respectively, and the large direct CP asymmetry (CPA) of 0.39 ± 0.17 for $B_s \rightarrow K^- \pi^+$ [5]. Moreover, surprisingly both detectors have also shown an unexpected large CP phase in the mixing-induced CPA for $B_s \rightarrow J/\Psi \phi$ with the average value given by [6]

$$\phi_s^{J/\Psi \phi} = -2\beta_s + \phi_s^{\text{NP}} = -2[0.10, 1.42] \quad (1)$$

at the 95% confidence level (CL), where $\beta_s \equiv \arg(-V_{ts}V_{tb}^*/V_{cs}V_{cb}^*) \approx 0.019$ [7] is the CP phase of the SM while ϕ_s^{NP} is a phase from new physics.

If the measurement of Eq. (1) indicates the existence of a new CP phase in $b \rightarrow s$ transition [7, 8], it is interesting to expose the effects in other $b \rightarrow s$ related phenomena, such as the puzzles of CPAs in $B \rightarrow K\pi$ and of polarizations in $B \rightarrow \phi K^*$. Nevertheless, as known that owing to the inevitable large uncertainty of nonperturbative QCD effects, it is difficult to definitely uncover the new CP phase from the CPAs of the nonleptonic exclusive decays. Therefore, the best environment to look for new phase is that QCD effects are less involved and SM contributions are highly suppressed. Therefore, we propose that the large CP violating phase in $b \rightarrow s$ transition can be directly probed by measuring T-odd observables in $\bar{B}_q \rightarrow V_q \ell^+ \ell^-$ decays with $V_q = K^*(\phi)$.

To understand the property of $\hat{T}$ violating approach for probing the new CP phase, we briefly discuss the characters of CP-odd and T-odd observables. In a decay process, the direct CPA or CP-odd observable is defined by $\mathcal{A}_{CP} \equiv (\bar{\Gamma} - \Gamma)/(\bar{\Gamma} + \Gamma)$, where Γ ($\bar{\Gamma}$) is the partial decay rate of the (CP-conjugate) process. Due to $\mathcal{A}_{CP} \propto \sin \theta_W \sin \theta_S$, it always needs weak CP phase θ_W and strong phase θ_S simultaneously to get the nonvanished CPA. Thus, the prediction efficiency on such CPA is mainly dictated by the uncertain calculations of strong phase. Another way to probe CP phase is through the spin-momentum triple correlation of $\vec{s} \cdot (\vec{p}_i \times \vec{p}_j)$ [9, 10], where $\vec{s}$ is the spin carried by one of outgoing particles and $\vec{p}_i$ and $\vec{p}_j$ denote any two independent momentum vectors. The triple momentum correlation is T-odd

observable since it changes sign under the time reversal ($\hat{T}$) transformation of $t \rightarrow -t$. We note that $\hat{T}$ is used to distinguish from real time-reversal transformation which also contains the interchange of initial and final states. By the $C\hat{P}\hat{T}$ invariant theorem, $\hat{T}$ violation (TV) implies CPV. Therefore, study of the T-odd observable can also help us to understand the origin of CPV. Intriguingly, the triple correlation is proportional to $\sin(\theta_W + \theta_S)$. This leads that the strong phase in T-odd observable is not necessary. To satisfy the criteria of a small QCD uncertainty with the high suppression in the SM for the $b \rightarrow s$ transition mentioned earlier, we find that TV in $B_q \rightarrow V_q \ell^+ \ell^-$ with polarized vector meson could provide a good chance to directly observe the new phase revealed by CDF and DØ.

To search for the new CP phase in the extension of the SM, a model independent analysis is not suitable for the current study. In this paper we investigate the minimal extension in the gauge sector by including one family dependent $U(1)'$ to the SM gauge symmetry, i.e. the considered gauge symmetry is $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)'_X$. Although lots of extensions of the SM predict additional $U(1)'$ gauge symmetry, such as string theory [11], grand unified theories [12], Little Higgs models [13], light U-boson model [16], etc, a well-motivated Z' model for low energy systems is the so-called nonuniversal Z' . Besides the $U(1)'$ charges carried by the fermions are family dependent [14], the model also leads to flavor changing neutral currents (FCNCs) at tree level and provides the solution to the puzzles in B physics [17]. The recent progress of the nonuniversal Z' on the phenomenological applications could be referred to Refs. [17, 18].

We start to set up the relevant interactions with the new Z' gauge particle. Following the convention in Refs. [14, 15], we write the couplings of the Z' -boson to fermions as

$$\mathcal{L}^{Z'} = -g_2 J_{Z'}^{f\mu} Z'_\mu, \quad (2)$$

with

$$J_{Z'}^{f\mu} = \bar{F} \gamma^\mu \left[\tilde{\epsilon}_f^L P_L + \tilde{\epsilon}_f^R P_R \right] F, \quad (3)$$

where $F^T = (f1, f2, f3)$ and the indexes fi ($i=1,2,3$) denote the same type of quarks (qs) or charged leptons (ℓs), $\tilde{\epsilon}_f^\chi$ are diagonal 3×3 matrices in flavor space with $\chi = L, R$ and are represented by $\text{diag} \tilde{\epsilon}_f^\chi = (X_{f1}^\chi, X_{f2}^\chi, X_{f3}^\chi)$. Note that X_{fi}^χ could be different for different generations in the nonuniversal Z' model. Hereafter, X_{fi}^χ are taken as free parameters. Although the third component of $SU(2)_L$ and $U(1)_Y$ in the SM also provide two neutral

gauge bosons, we will not show them explicitly since they involve only flavor conserved processes and are well known. To obtain the interactions in physical states, we introduce V_f^χ to diagonalize the associated Yukawa matrix of fermion. In terms of mass eigenstates, Eq. (3) could be written as

$$\begin{aligned} J_{Z'}^{f\mu} &= \bar{F}\gamma^\mu [\mathbf{L}^f P_L + \mathbf{R}^f P_R] F, \\ \mathbf{L}^f &= V_f^L \tilde{\boldsymbol{\epsilon}}_f^L V_f^{L\dagger}, \quad \mathbf{R}^f = V_f^R \tilde{\boldsymbol{\epsilon}}_f^R V_f^{R\dagger}. \end{aligned} \quad (4)$$

Due to the unequal elements in $\tilde{\boldsymbol{\epsilon}}_f^\chi$, obviously, flavor misaligned effects in $\mathbf{L}^f$ and $\mathbf{R}^f$ will lead to FCNCs at tree level.

In order to display the new physics effects with more compact form in the processes, we use vector and axial-vector currents instead chiral currents. Thus, Eq. (4) is rewritten by

$$\begin{aligned} J_{Z'}^{f\mu} &= \bar{F}\gamma^\mu [\mathbf{c}_V^f + \mathbf{c}_A^f \gamma_5] F, \\ \mathbf{c}_V^f &= \mathbf{R}^f + \mathbf{L}^f, \quad \mathbf{c}_A^f = \mathbf{R}^f - \mathbf{L}^f. \end{aligned} \quad (5)$$

Consequently, the Z' -mediated interactions for $b \rightarrow s$ are given by

$$\mathcal{L}_{b \rightarrow s}^{Z'} = -g_2 \bar{s} \gamma^\mu (c_{Vsb}^q + c_{Asb}^q \gamma_5) b Z'_\mu + h.c. \quad (6)$$

For semileptonic decays, the couplings to the leptons are taken as

$$\mathcal{L}_{\ell\ell}^{Z'} = -g_2 \bar{\ell} \gamma^\mu (c_{V\ell\ell}^\ell + c_{A\ell\ell}^\ell \gamma_5) \ell Z'_\mu. \quad (7)$$

Since we will concentrate on the $\ell^+ \ell^-$ -pair production, to avoid the problem on the lepton flavor violation and nonuniversality and simplify our analysis, we assume that $c_{V(A)\ell\ell}^\ell$ are the same for different families, i.e. $c_{V(A)ee}^\ell = c_{V(A)\mu\mu}^\ell = c_{V(A)\tau\tau}^\ell$. In the following analysis, we will show that the impacts of Z' -mediated FCNCs on the processes of B_s mixing, $\bar{B}_s \rightarrow \mu^+ \mu^-$ and $\bar{B}_q \rightarrow V_q \ell^+ \ell^-$. Note that since we don't have any information on V_q^χ except the CKM matrix defined by $\mathbf{V} = V_U^L V_D^{L\dagger}$, the elements in $\mathbf{c}_{V,A}^q$ should be regarded as free parameters. Therefore, the parameters associated with the processes for $b \rightarrow d$ and $s \rightarrow d$ transitions could be independent of those associated with the $b \rightarrow s$ transition.

Based on the new effects, we now formulate the related phenomena for $\Delta B = 2$ and $\Delta B = 1$ processes. In terms of Eq. (6), the tree induced effective Hamiltonian for $\Delta B = 2$ is read by

$$\mathcal{H}_{\Delta B=2}^{Z'} = \frac{G_F \omega}{2\sqrt{2}} [\bar{s} \gamma_\mu (c_{Vsb}^q + c_{Asb}^q \gamma_5) b]^2, \quad (8)$$

where $G_F/\sqrt{2} = g_1^2/8m_W^2$, g_1 is the gauge coupling of $SU(2)_L$ and $\omega = 4m_W^2(g_2/g_1)^2/m_{Z'}^2$. Including the Wick contracting factor and using the matrix elements parametrized by

$$\begin{aligned}\langle \bar{B} | \bar{s} \gamma_\mu P_\chi b \bar{s} \gamma^\mu P_\chi b | B \rangle &= \frac{1}{3} m_B f_B^2 \hat{B}_B, \\ \langle \bar{B} | \bar{s} \gamma_\mu P_L b \bar{s} \gamma^\mu P_R b | B \rangle &= -\frac{5}{12} m_B f_B^2 \hat{B}_B,\end{aligned}\tag{9}$$

we obtain

$$\begin{aligned}M_{12}^{Z'} &= \langle \bar{B} | \mathcal{H}_{\Delta B=2}^{Z'} | B \rangle, \\ &= \frac{G_F \omega}{24\sqrt{2}} m_B \hat{B}_B f_B^2 [-(c_{Vsb}^q)^2 + 9(c_{Asb}^q)^2].\end{aligned}\tag{10}$$

Combining the result with the SM contribution, the transition matrix element for $\bar{B}_s \rightarrow B_s$ is given by

$$M_{12}^s = A_{12}^{\text{SM}} e^{-2i\beta_s} + A_{12}^{Z'} e^{2i(\theta_s^{Z'} - \beta_s)},\tag{11}$$

where $A_{12}^{Z'} = |M_{12}^{Z'}|$ and the CP phase of $M_{12}^{Z'}$ is parametrized by $2(\theta_s^{Z'} - \beta_s)$ [7]. As a result, the B_s mixing is obtained as

$$\Delta m_s = 2|M_{12}^s| = \Delta m_s^{\text{SM}} \left(1 + 2r \cos 2\theta_s^{Z'} + |r|^2\right)^{1/2}\tag{12}$$

with $r = A_{12}^{Z'}/A_{12}^{\text{SM}}$. Due to $\Delta\Gamma_s \ll \Delta m_s$ in the B-system, the time-dependent CPA could be simplified to be

$$\begin{aligned}-S_{J/\Psi\phi} &\simeq \text{Im} \left(\sqrt{\frac{M_{12}^{s*}}{M_{12}^s}} \right) = \sin(2\beta_s - \phi_s^{Z'}), \\ \phi_s^{Z'} &= \arctan \left(\frac{r \sin 2\theta_s^{Z'}}{1 - r \cos 2\theta_s^{Z'}} \right).\end{aligned}\tag{13}$$

For $\Delta B = 1$ processes, we focus on the important decay mode of $b \rightarrow s \ell^+ \ell^-$. From Eqs. (6) and (7), the corresponding effective interactions can be found to be

$$\mathcal{H}_{b \rightarrow s \ell \ell}^{Z'} = \frac{G_F \omega}{2\sqrt{2}} \bar{s} \gamma_\mu (c_{Vsb}^q + c_{Asb}^q \gamma_5) b \bar{\ell} \gamma^\mu (c_{V\ell\ell}^\ell + c_{A\ell\ell}^\ell \gamma_5) \ell.\tag{14}$$

The direct application of Eq. (14) is the chiral suppressed decay of $\bar{B}_s \rightarrow \ell^+ \ell^-$. Using $\langle 0 | \bar{s} \gamma_\mu \gamma_5 b | \bar{B}_s(p) \rangle = i f_{B_s} p_\mu$ and including the SM contributions, the branching ratio (BR) for $\bar{B}_s \rightarrow \ell^+ \ell^-$ is expressed by

$$\mathcal{B}(\bar{B}_s \rightarrow \ell^+ \ell^-) = \tau_{B_s} \frac{G_F^2}{16\pi} m_{B_s} f_{B_s}^2 m_\ell^2 \left(1 - \frac{4m_\ell^2}{m_{B_s}^2}\right)^{1/2} |A_{\ell\ell}|^2$$

with

$$\begin{aligned} A_{\ell\ell} &= \frac{\lambda_t \alpha}{\pi \sin^2 \theta_W} Y(x_t) + \omega c_{Asb}^q c_{Al}^\ell, \\ \omega &= \left(\frac{g_2}{g_1} \right)^2 \frac{4m_W^2}{m_{Z'}^2}, \end{aligned} \quad (15)$$

where $\lambda_t = V_{ts}^* V_{tb}$, θ_W is the Weinberg angle, $\alpha = e^2/4\pi$ and $Y(x_t) \approx 0.315x_t^{0.78}$ with $x_t = m_t^2/m_W^2$ [21].

To study the $\hat{T}$ violating effects in $\bar{B}_q \rightarrow V_q \ell^+ \ell^-$ decays, besides the new interactions of Eq. (14), we need the hadronic effects for the $\bar{B}_q \rightarrow V_q$ transition. As usual, the form factors associated with various current operators are parametrized as

$$\begin{aligned} \langle V(p_V, \epsilon) | V_\mu | \bar{B}(p_B) \rangle &= i \frac{V(q^2)}{m_B + m_V} \varepsilon_{\mu\alpha\beta\rho} \epsilon^{*\alpha} P^\beta q^\rho, \\ \langle V(p_V, \epsilon) | A_\mu | \bar{B}(p_B) \rangle &= 2m_V A_0(q^2) \frac{\epsilon^* \cdot q}{q^2} q_\mu \\ &\quad + (m_B + m_V) A_1(q^2) \left(\epsilon_\mu^* - \frac{\epsilon^* \cdot q}{q^2} q_\mu \right) \\ &\quad - A_2(q^2) \frac{\epsilon^* \cdot q}{m_{B_q} + m_{V_q}} \left(P_\mu - \frac{P \cdot q}{q^2} q_\mu \right), \\ \langle V(p_V, \epsilon) | T_{\mu\nu} q^\nu | \bar{B}(p_B) \rangle &= -iT_1(q^2) \varepsilon_{\mu\alpha\beta\rho} \epsilon^{*\alpha} P^\beta q^\rho, \\ \langle V(p_V, \epsilon) | T_{\mu\nu}^5 q^\nu | \bar{B}(p_B) \rangle &= T_2(q^2) \left(\epsilon_\mu^* P \cdot q - \epsilon^* \cdot q P_\mu \right) \\ &\quad + T_3(q^2) \epsilon^* \cdot q \left(q_\mu - \frac{q^2}{P \cdot q} P_\mu \right) \end{aligned} \quad (16)$$

with $V_\mu = \bar{s} \gamma_\mu b$, $A_\mu = \bar{s} \gamma_\mu \gamma_5 b$, $T_{\mu\nu} = \bar{s} i \sigma_{\mu\nu} b$, $T_{\mu\nu}^5 = \bar{s} i \sigma_{\mu\nu} \gamma_5 b$, $P = p_B + p_V$ and $q = p_B - p_V$. Here, we have used B (V) instead of B_q (V_q) and $m_B(m_V) = m_{B_q}(m_{V_q})$. Hereafter, we will adopt the notation for shorthand. Hence, the transition amplitude for $\bar{B} \rightarrow V \ell^+ \ell^-$ ($\ell = e, \mu$) with the polarized V can be arranged by [10]

$$\begin{aligned} \mathcal{M}_V^{(\lambda)} &= -\frac{G_F \alpha \lambda_t}{2\sqrt{2}\pi} \left\{ \mathcal{M}_{1\mu}^{(\lambda)} L^\mu + \mathcal{M}_{2\mu}^{(\lambda)} L^{5\mu} \right\}, \\ \mathcal{M}_{a\mu}^{(\lambda)} &= i\xi_1 \varepsilon_{\mu\nu\alpha\beta} \epsilon^{*\nu}(\lambda) P^\alpha q^\beta + \xi_2 \epsilon_\mu^*(\lambda) + \xi_3 \epsilon^* \cdot q P_\mu, \end{aligned}$$

where $L_\mu = \bar{\ell} \gamma_\mu \ell$, $L_\mu^5 = \bar{\ell} \gamma_\mu \gamma_5 \ell$, subscript $a = 1[2]$ while $\xi_i = h_i [g_i]$ ($i = 1, 2, 3$) and their

explicit expressions are

$$\begin{aligned}
h_1 &= [C_9^{\text{eff}} + c_{Vsb}^q c_{V\ell\ell}^\ell \zeta] \frac{V}{m_B + m_V} + \frac{2m_b}{q^2} C_7 T_1, \\
h_2 &= [-C_9^{\text{eff}} + c_{Asb}^q c_{V\ell\ell}^\ell \zeta] (m_B + m_V) A_1 P \cdot q C_7 T_2, \\
h_3 &= [C_9^{\text{eff}} - c_{Asb}^q c_{V\ell\ell}^\ell \zeta] \frac{A_2}{m_B + m_V} + \frac{2m_b}{q^2} C_7 \left(T_2 + \frac{q^2}{P \cdot q} T_3 \right), \\
g_1 &= [C_{10} + c_{Vsb}^q c_{A\ell\ell}^\ell \zeta] \frac{V}{m_B + m_V}, \\
g_2 &= [-C_{10} + c_{Asb}^q c_{A\ell\ell}^\ell \zeta] (m_B + m_V) A_1, \\
g_3 &= [C_{10} - c_{Asb}^q c_{A\ell\ell}^\ell \zeta] \frac{A_2}{m_B + m_V}.
\end{aligned} \tag{17}$$

Here, we have suppressed the q^2 -dependence in the form factors, $(C_9^{\text{eff}}, C_7, C_{10})$ are the Wilson coefficients of the SM and their specific results could be referred to Ref. [21] and $\zeta = \pi\omega/\alpha\lambda_t$.

To obtain the $\hat{T}$ violating observables, which are connected to the polarizations of the vector meson in $\bar{B} \rightarrow V\ell^+\ell^-$ decays, we have to consider the decay chain $\bar{B} \rightarrow V(\rightarrow P_1 P_2)\ell^+\ell^-$ in which $P_1 P_2$ is $K\pi(KK)$ as $V = K^*(\phi)$. For calculations, we set the positron lepton momentum as $p_{\ell^+} = \sqrt{q^2}/2(1, \sin\theta_{\ell^+}, 0, \cos\theta_{\ell^+})$ in the q^2 rest frame and the K momentum $p_K = m_V/2(1, \sin\theta_K \cos\phi_K, \sin\theta_K \sin\phi_K, \cos\theta_K)$ in the V rest frame. With some tedious algebra, the differential decay rate with polarized V-meson can be formulated by [10, 22]

$$\begin{aligned}
\frac{d\Gamma}{d\Omega_K dq^2} &= \frac{G_F^2 \alpha^2 |\lambda_t|^2 |\vec{p}|}{2^{14} \pi^6 m_B^2} \mathcal{B}(K^*[\phi] \rightarrow K\pi[KK]) \\
&\times \left\{ 16 \cos^2 \theta_K \sum_{a=1,2} |\mathcal{M}_a^0|^2 + 8 \sin^2 \theta_K \sum_{a=1,2} \left(|\mathcal{M}_a^+|^2 + |\mathcal{M}_a^-|^2 \right) \right. \\
&- 8 \sin^2 \theta_K \left[\cos 2\phi_K \sum_{a=1,2} \text{Re} \mathcal{M}_a^+ \mathcal{M}_a^{-*} + \sin 2\phi_K \sum_{a=1,2} \text{Im} \mathcal{M}_a^+ \mathcal{M}_a^{-*} \right] \\
&+ 3\pi \sin 2\theta_K \left[\cos \phi_K \left(\text{Re} \mathcal{M}_1^0 (\mathcal{M}_2^{+*} - \mathcal{M}_2^{-*}) + \text{Re} (\mathcal{M}_1^+ - \mathcal{M}_1^-) \mathcal{M}_2^{0*} \right) \right. \\
&\left. \left. + \sin \phi_K \left(\text{Im} \mathcal{M}_1^0 (\mathcal{M}_2^{+*} + \mathcal{M}_2^{-*}) - \text{Im} (\mathcal{M}_1^+ + \mathcal{M}_1^-) \mathcal{M}_2^{0*} \right) \right] \right\}
\end{aligned} \tag{18}$$

where $d\Omega_K = d\cos\theta_K d\phi_K$, θ_K is the polar angle of K-meson while the ϕ_K is the relative angle of decaying planes of $P_1 P_2$ and $\ell^+\ell^-$, $|\vec{p}| = [((m_B^2 + m_V^2 - q^2)/(2m_B))^2 - m_V^2]^{1/2}$, $\mathcal{M}_a^0$ and $\mathcal{M}_a^\pm$ denote the longitudinal and transverse polarizations of V respectively and they are

given as follows:

$$\begin{aligned}\mathcal{M}_a^0 &= \sqrt{q^2} \left(\frac{E_V}{m_V} \xi_2 + 2\sqrt{q^2} \frac{|\vec{p}_V|^2}{m_V} \xi_3 \right), \\ \mathcal{M}_a^\pm &= \sqrt{q^2} \left(\pm 2|\vec{p}_V| \sqrt{q^2} \xi_1 + \xi_2 \right)\end{aligned}$$

with $|\vec{p}_V| = \sqrt{E_V^2 - m_V^2}$ and $E_V = (m_B^2 - m_V^2 - q^2)/2\sqrt{q^2}$. Since we have integrated out the angles associated with the lepton, some important effect such as the forward-backward asymmetry (FBA) of the lepton does not show up in Eq. (18). However, we find an up-down (UD) asymmetry of the K -meson, which is associated with $\sin 2\theta_K$ and similar to the lepton FBA, could be generated.

According to Eq. (18), it is clear that besides the real pieces, the imaginary parts that represent the TV (or CPV) also appear in the angular distribution. If we can find the proper T -odd operators denoted by $\mathcal{O}_{T_i}$, in terms of the concept of quantum mechanics, the average of the T -odd operator can be written as

$$\langle \mathcal{O}_{T_i} \rangle = \int d\Omega_K \mathcal{O}_{T_i} \text{sign}(\omega_i) \frac{d\Gamma}{d\Omega_K dq^2}$$

where $\text{sign}(\omega_i)$ is chosen to get a nonvanished asymmetric observables when integrating the angles of phase space. Then, the T -odd observable could be defined by

$$\mathcal{A}_{T_i} = \frac{\langle \mathcal{O}_{T_i} \rangle}{\int d\Omega_K (d\Gamma/d\Omega_K dq^2)}.$$

Accordingly, we find that the two $\hat{T}$ violating terms in Eq. (18) can be extracted by using T -odd momentum correlations, defined by

$$\begin{aligned}\mathcal{O}_{T_1} &= 4|\vec{p}_B| \frac{(\vec{p}_B \cdot \vec{p}_{\ell^+} \times \vec{p}_K)(\vec{p}_B \times \vec{p}_K) \cdot (\vec{p}_{\ell^+} \times \vec{p}_B)}{|\vec{p}_B \times \vec{p}_K|^2 |\vec{p}_{\ell^+} \times \vec{p}_B|^2}, \\ &= 2 \sin 2\phi_K, \\ \mathcal{O}_{T_2} &= \frac{16}{3\pi} \frac{|\vec{p}_B| |\vec{p}_K| \cdot (\vec{p}_B \times \vec{p}_{\ell^+})}{|\vec{p}_B \times \vec{p}_K| |\vec{p}_B \times \vec{p}_{\ell^+}|} = \frac{16}{3\pi} \sin \phi_K\end{aligned}$$

where the momenta of particles are taken in the V -meson rest frame, $\text{sign}(\omega_1) = \text{sign}(\sin \theta_K)$ and $\text{sign}(\omega_2) = \text{sign}(\cos \theta_K)$. As a result, we get

$$\mathcal{A}_{T_1} = - \frac{\sum_{a=1,2} \text{Im}(\mathcal{M}_a^+ \mathcal{M}_a^{-*})}{\sum_{\lambda=+,0,-} \sum_{a=1,2} |\mathcal{M}_a^\lambda|^2}, \quad (19)$$

$$\mathcal{A}_{T_2} = \frac{1}{\sum_{\lambda=+,0,-} \sum_{a=1,2} |\mathcal{M}_a^\lambda|^2} \left[\text{Im} \mathcal{M}_1^0 (\mathcal{M}_2^{+*} + \mathcal{M}_2^{-*}) - \text{Im}(\mathcal{M}_1^+ + \mathcal{M}_1^-) \mathcal{M}_2^{0*} \right]. \quad (20)$$

Although our main purpose is to explore the existence of new CP phase, the Z' -mediated effects also have significant contributions to CP-conserved quantities. Therefore, besides the physical observables defined in Eqs. (19) and (20), we could also invent interesting measurable quantities, such as longitudinal and transverse polarizations of the V -meson and the angular asymmetry of the K -meson, etc [10]. According to the definition in Ref. [10], the longitudinal and transverse polarizations of V -meson appearing in Eq. (18) can be found as

$$\mathcal{P}_L(q^2) = \frac{\sum_{a=1,2} |\mathcal{M}_a^0|^2}{\sum_{\lambda=+,0,-} \sum_{a=1,2} |\mathcal{M}_a^\lambda|^2}, \quad (21)$$

$$\mathcal{P}_T(q^2) = \frac{\sum_{a=1,2} |\mathcal{M}_a^+|^2 + |\mathcal{M}_a^-|^2}{\sum_{\lambda=+,0,-} \sum_{a=1,2} |\mathcal{M}_a^\lambda|^2}. \quad (22)$$

In addition, due to the appearance of $\sin 2\theta_K \cos \phi_K$ in Eq. (18), the UD asymmetry of the K -meson can be obtained as [10]

$$\mathcal{A}_{UD}^K = \frac{1}{\sum_{\lambda=+,0,-} \sum_{a=1,2} |\mathcal{M}_a^\lambda|^2} \left[\text{Re} \mathcal{M}_1^0 (\mathcal{M}_2^{+*} - \mathcal{M}_2^{-*}) + \text{Re} (\mathcal{M}_1^+ - \mathcal{M}_1^-) \mathcal{M}_2^{0*} \right]. \quad (23)$$

So far we have introduced seven independent free parameters of ω , $|c_{Vsb}^q|e^{i\theta_V}$ and $|c_{Asb}^q|e^{i\theta_A}$ and $c_{V\ell\ell}^\ell$ and $c_{A\ell\ell}^\ell$ in our analysis. In order to further reduce the number of free parameters to simplify the phenomenological analysis, we set $g_1 = g_2$ and $m_{Z'} = 1\text{TeV}$ and force $\theta_V = \theta_A$. Although the restricted condition of the latter is unnecessary, it could correspond to a special scenario of Yukawa matrices, e.g. hermitian Yukawa matrices which result in $V_f^L = V_f^R$ [23]. For numerical estimations, we need the data to constrain the remaining five parameters. Therefore, we adopt the values of $\Delta m_s = 18.17 \pm 0.86 \text{ ps}^{-1}$ [3, 4], $\mathcal{B}(\bar{B}_d \rightarrow K^{[*]0} \mu^+ \mu^-) = (0.45_{-0.10}^{+0.12}) [1.05_{-0.13}^{+0.15}] \times 10^{-6}$ [2] and $\mathcal{B}(\bar{B}_s \rightarrow \mu^+ \mu^-) < 4.7 \times 10^{-8}$ [26] as the inputs whereas the SM predictions are taken as $\Delta m_s^{\text{SM}} = 17.37 \pm 6.03 \text{ ps}^{-1}$ [24], $\mathcal{B}(\bar{B}_d \rightarrow K^{[*]0} \mu^+ \mu^-)_{\text{SM}} = 0.58 [1.3] \times 10^{-6}$ [25] and $\mathcal{B}(\bar{B}_s \rightarrow \mu^+ \mu^-)_{\text{SM}} \approx 0.33 \times 10^{-8}$ [27]. Since $\bar{B}_s \rightarrow \phi \ell^+ \ell^-$ is very similar to $\bar{B} \rightarrow K^* \ell^+ \ell^-$, we use the latter as the delegate.

We now present the numerical results. In order to realize the strong correlation among $\phi_s^{J/\Psi\phi}$, Δm_s , $\mathcal{B}(\bar{B}_s \rightarrow \mu^+ \mu^-)$ and $\mathcal{B}(\bar{B}_d \rightarrow K^{[*]0} \ell^+ \ell^-)$ influenced by the same parameters, we show the observables versus Δm_s in Fig. 1. Here, for the $\bar{B} \rightarrow K^{(*)}$ transition form factors, we employ the results calculated by light-cone sum rules (LCSRs) [28]. By the figure, we see clearly that besides the BR of $\bar{B}_s \rightarrow \mu^+ \mu^-$ can be enhanced to be close to $O(10^{-8})$, the large $\phi_s^{J/\Psi\phi}$ generated by Z' -mediated effects can be consistent with the current measurements of




FIG. 1: Correlations between $(\phi_s^{J/\Psi\phi}, \mathcal{B}(\bar{B}_s \rightarrow \mu^+\mu^-), \mathcal{B}(\bar{B}_d \rightarrow K^{*0}\mu^+\mu^-), \mathcal{B}(\bar{B}_d \rightarrow K^0\mu^+\mu^-))$ and Δm_s by the influence of nonuniversal Z' .

DØ and CDF. With the values of parameters which are constrained by the current data of Δm_s , $\mathcal{B}(\bar{B}_s \rightarrow \mu^+\mu^-)$ and $\mathcal{B}(\bar{B}_d \rightarrow K^{[*]0}\ell^+\ell^-)$, the predicted T-odd observables of $\mathcal{A}_{T_{1,2}}$ are displayed in Fig. 2. It is clear that we can obtain $\mathcal{A}_{T_1}$ of 10% and $\mathcal{A}_{T_2}$ of 20% at large and small q^2 , respectively. Moreover, with the same set of values of parameters, the predicted




FIG. 2: T-odd observables $\mathcal{A}_{T_{1,2}}$ in $\bar{B}_d \rightarrow K^{*0}\mu^+\mu^-$ as a function of q^2 in nonuniversal Z' model.

longitudinal [transverse] polarization and the UD asymmetry defined in Eq. (21) [Eq. (22)] and Eq. (23) are given in Fig. 3(a) and (b), respectively, where the solid and dashed lines denote the SM predictions. We find that the deviation of 10% from the SM value in $\mathcal{P}_{L[T]}$ can be achieved in the Z' models. Additionally, the crossing of $\mathcal{P}_L$ and $\mathcal{P}_T$ at large q^2 is

also sensitive to the new physics effects. Besides a large deviation from the SM result in the spectrum of $\mathcal{A}_{UD}^K$ could be accomplished, the zero point of $\mathcal{A}_{UD}^K$ is also sensitive to the Z' -mediated effects.




FIG. 3: (a) Longitudinal [Transverse] polarization $\mathcal{P}_{L[T]}$ of K^* and (b) up-down asymmetry of K in $\bar{B}_d \rightarrow K^{*0}(\rightarrow K\pi)\mu^+\mu^-$, where solid and dashed lines stand for the SM predictions.

In summary, we have studied the Z' -mediated effects on Δm_s , $\phi_s^{J/\Psi\phi}$, $\bar{B}_s \rightarrow \mu^+\mu^-$ and $\bar{B} \rightarrow V\ell^+\ell^-$ in the family dependent $U(1)'$ model. With the constraints of the current data, we have found that in the nonuniversal Z' model, the time-dependent CP violating phase of the B_s oscillation could be $\phi_s^{J/\Psi\phi} = -(0.26 \pm 0.22)$ and the T-odd observables $\mathcal{A}_{T_1}$ and $\mathcal{A}_{T_2}$ could be 10% and 20% at the large and small q^2 regions, respectively. In addition, we have also shown that the longitudinal and transverse polarizations of K^* and the UD asymmetry of K in the decay chain $\bar{B} \rightarrow K^*(\rightarrow K\pi)\ell^+\ell^-$ could be significantly influenced by the Z' effects.

Acknowledgments

This work is supported in part by the National Science Council of R.O.C. under Grant No: NSC-97-2112-M-006-001-MY3.

[1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).

- [2] E. Barberio *et al.*, arXiv:0808.1297 [hep-ex], online update at <http://www.slac.stanford.edu/xorg/hfag>.
- [3] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. Lett. **97**, 242003 (2006) [arXiv:hep-ex/0609040].
- [4] DØ note 5474-conf at <http://www-d0.fnal.gov/Run2Physics/WWW/results/prelim/B/B51>.
- [5] M. Morello (CDF Collaboration), Phys. Conf. Ser. **110**, 052040 (2008) [arXiv:0711.0693 [hep-ex]].
- [6] DØ note 5928-conf at <http://www-d0.fnal.gov/Run2Physics/WWW/results/prelim/B/B59>.
- [7] C. H. Chen, C. Q. Geng and L. Li, Phys. Lett. **B670**, 374 (2009) [arXiv:0808.0127 [hep-ph]].
- [8] F. J. Botella, G. C. Branco and M. Nebot, Phys. Rev. D **79**, 096009 (2009) [arXiv:0805.3995 [hep-ph]].
- [9] C. H. Chen and C. Q. Geng, Phys. Rev. D **64**, 074001 (2001) [arXiv:hep-ph/0106193].
- [10] C. H. Chen and C. Q. Geng, Nucl. Phys. **B636**, 338 (2002) [arXiv:hep-ph/0203003]; Phys. Rev. D **66**, 014007 (2002) [arXiv:hep-ph/0205306]; Phys. Rev. D **66**, 094018 (2002) [arXiv:hep-ph/0209352].
- [11] M. Cvetič and P. Langacker, Phys. Rev. D **54**, 3570 (1996) [arXiv:hep-ph/9511378].
- [12] R. W. Robinett and J. L. Rosner, Phys. Rev. D **25**, 3036 (1982) [Erratum *ibid.* D **27**, 679 (1983)]; *ibid.* **26**, 2396 (1982).
- [13] N. Arkani-Hamed, A. G. Cohen and H. Georgi, Phys. Lett. **B513**, 232 (2001) [arXiv:hep-ph/0105239].
- [14] P. Langacker and M. Plümacher, Phys. Rev. D **62**, 013006 (2000) [arXiv:hep-ph/0001204].
- [15] P. Langacker and M. Luo, Phys. Rev. D **45**, 278 (1992).
- [16] P. Fayet, Phys. Rev. D **75**, 115017 (2007) [arXiv:hep-ph/0702176]; C. H. Chen, C. Q. Geng and C. W. Kao, Phys. Lett. **B663**, 400 (2008) [arXiv:0708.0937 [hep-ph]].
- [17] V. Barger *et al.*, Phys. Lett. **B580**, 186 (2004) [arXiv:hep-ph/0310073]; V. Barger *et al.*, Phys. Lett. **B596**, 229 (2004) [arXiv:hep-ph/0405108]; V. Barger *et al.*, Phys. Lett. **B598**, 218 (2004) [arXiv:hep-ph/0406126]; K. Cheung *et al.*, Phys. Lett. **B652**, 285 (2007) [arXiv:hep-ph/0604223]; V. Barger *et al.*, Phys. Rev. D **80**, 055008 (2009) [arXiv:0902.4507 [hep-ph]]; arXiv:0906.3745.
- [18] C. H. Chen and H. Hatanaka, Phys. Rev. D **73**, 075003 (2006) [arXiv:hep-ph/0602140]; Q. Chang, X. Q. Li and Y. D. Yang, JHEP **0905**, 056 (2009) [arXiv:0903.0275 [hep-ph]].

- X. G. He and G. Valencia, Phys. Lett. **B680**, 72 (2009) [arXiv:0907.4034 [hep-ph]]; C. W. Chiang *et al.*, arXiv:0910.2929 [hep-ph]; C. W. Chiang, R. H. Li and C. D. Lu, arXiv:0911.2399 [hep-ph].
- [19] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **100**, 161802 (2008) [arXiv:0712.2937 [hep-ex]].
- [20] V. M. Abazov *et al.* (DØ Collaboration), arXiv:0802.2255 [hep-ex].
- [21] G. Buchalla, A.J. Buras and M.E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996) [arXiv:hep-ph/9512380].
- [22] C. Bobeth, G. Hiller and G. Piranishvili, JHEP **0807**, 106 (2008) [arXiv:0805.2525 [hep-ph]]; U. Egede *et al.*, JHEP **0811**, 032 (2008) [arXiv:0807.2589 [hep-ph]].
- [23] R. N. Mohapatra and G. Senjanovic, Phys. Lett. **B79**, 283 (1978); C. H. Chen, Phys. Lett. **B521**, 315 (2001) [arXiv:hep-ph/0110098]. .
- [24] A. Lenz and U. Nierste, JHEP **06**, 072 (2007) [arXiv:hep-ph/0612167]; A. Lenz, arXiv:0802.0977 [hep-ph].
- [25] C. H. Chen and C. Q. Geng, Mod. Phys. Lett. **A21**, 1137 (2006) [arXiv:hep-ph/0502246].
- [26] T. Aaltonen *et al.* (CDF Collabpration), Phys. Rev. Lett. **100**, 101802 (2008) [arXiv:0712.1708 [hep-ex]].
- [27] M. Blanke *et al.*, JHEP **0610**, 003 (2006) [arXiv:hep-ph/0604057].
- [28] P. Ball and R. Zwicky, Phys. Rev. **D71**, 014015 (2005) [arXiv:hep-ph/0406232]; *ibid*, 014029 (2005) [arXiv:hep-ph/0412079].